

## Geographic distribution of Apicomplexa infecting *Strombus gigas*

DALILA ALDANA ARANDA<sup>1</sup>, LILIANE FRENKIEL<sup>2</sup>, ERICK BAQUEIRO CÁRDENAS<sup>3</sup>, ADRIANA ZETINA ZÁRATE<sup>1</sup>, GRACIELA GARCÍA MOLINER<sup>4</sup>, AURY RODRÍGUEZ<sup>5</sup>, JEANNETTE MATEO PÉREZ<sup>6</sup>, ALEJANDRO TAGLIAFICO<sup>7</sup>, ERICK CASTRO<sup>8</sup>, TOMÁS CAMARENA<sup>9</sup> AND GUSTAVO ARENCIBIA<sup>10</sup>

<sup>1</sup>CINVESTAV IPN Unidad Mérida, laboratorio de Biología y Cultivo de Moluscos

<sup>2</sup>Université des Antilles et de la Guyane, French West Indies, <sup>3</sup>CICATA IPN Altamira, México; <sup>4</sup>CFMC, Puerto Rico,

<sup>5</sup>Departamento de Ciencias Marinas, Universidad de Puerto Rico -Mayaguez

<sup>6</sup>Dirección de Recursos Pesqueros, Secretaría del Medio Ambiente, República Dominicana

<sup>7</sup>Universidad de Oriente, Núcleo Nueva Esparta, Venezuela; <sup>8</sup>Secretaría de Agricultura y Pesca, San Andrés, Colombia

<sup>9</sup>Sistema Arrecifal Mesoamericano, Belice; <sup>10</sup>Centro de Investigaciones Pesqueras, Cuba

### ABSTRACT

The queen conch *Strombus gigas* is a marine resource of ecological and economical importance in the Caribbean region that suffers a reduction of its populations in most Caribbean countries. An intense and generalized sporozoan infection was detected in *Strombus gigas* populations from various Caribbean countries. The parasite, apparently a Coccidian, Apicomplexa was observed on histological sections of the digestive gland of every sampled organism throughout the various sites: in the samples from San Andres, Colombia; Guadeloupe and St. Barthelemy, French West Indies; Alacranes reef and Chinchorro bank, Mexico; Parguera, Puerto Rico; Margarita Island, Venezuela. To the present light-microscopy analysis, the parasites from all localities could be the same and apparently, they complete their cycle in the digestive gland tissue. The parasite was abundant in Puerto Rico, Colombia and Mexico samples, coinciding with observed anomalies in reproductive cycle, reduced gametogenesis and maturity in Alacranes, Mexico (20-40%) and no gonad activity was observed in conchs from San Andrés. The damage to the digestive diverticula begins in the cryptic cells and attains secretory cells later, giving way to discharge of cysts to the stomach through the digestive gland ducts. Various stages are identified. Five different stages were identified: Apicomplexa-like trophozoites, gametocysts, gamonts, gametes and oocysts. Given the presence of multiple stages at any time a re-infection and the life cycle is assumed to occur within the same host. These results raise several questions: Given the generalized infection at so distant sites, if the parasite is the same, are *S. gigas* populations more connected than it has been supposed? What are the environmental factors inducing such an intense and generalized infection? What is the impact on recruitment and therefore on the fishery? It is necessary to identify the geographic distribution of this parasite, to discriminate damages to the digestive gland from the physiological cycle destruction of digestive gland cells; to evaluate its impact on the reproductive activity.

KEY WORDS: *Strombus gigas*, parasites, queen conch, Caribbean

### Distribución Geográfica de Apicomplexa Infectando a *Strombus gigas*

El caracol *Strombus gigas* es un recurso marino con importancia económica y ecológica en el Caribe. Una intensa y generalizada infección de un esporozoario presentan varias poblaciones de *Strombus gigas* en varias localidades del Caribe: Turmaline, Puerto Rico; San Andrés, Colombia; Guadalupe y St. Barthelemy, Antillas Francesas; arrecife Alacranes y banco Chinchorro, México; Isla Margarita, Venezuela. El parásito es probablemente un Apicomplexa encontrado en la glándula digestiva de *S. gigas* en diferentes localidades, presentándose una invasión frecuente del tejido alveolar de la glándula digestiva. El daño de los divertículos digestivos incluye tanto a las células crípticas como las secretoras. A nivel microscópico el parásito parece ser el mismo en las localidades estudiadas y pareciera completar su ciclo en el mismo organismo. La infección fue abundante en los organismos de Puerto Rico, Colombia y México, coincidiendo con anomalías en el ciclo reproductor. Los organismos de Alacranes presentaron una reducción de la gametogénesis y madurez (20-40%), mientras que los de San Andrés, no presentaron actividad gonádica. El daño en la glándula digestiva se inicia en las células crípticas, invadiendo posteriormente las células secretoras, liberando quistes al estómago a través de los ductos digestivos. Se identificaron varios estadios: Trofozoito tipo Apicomplexa, gametocistos, gamontes, gametos y oocistos. Dada la presencia de los diferentes estadios en un mismo organismo, se presupone una posible re-infección y el ciclo completo en el hospedero. Los resultados anteriores generan varias preguntas: ¿Dada la infección generalizada, eso implicaría que las poblaciones de *S. gigas* están más interconectadas que lo que se supone? Cuáles son las condiciones ambientales que favorecen esta infección?Cuál es su impacto en el reclutamiento?

PALABRAS CLAVES: *Strombus gigas*, parásitos, caracol rosa, Caribe

## INTRODUCTION

Given the regional importance of *Strombus gigas* Linne, 1758 in the Caribbean, and the critical state of some of its populations (Appeldoorn, 1987; Aldana Aranda et al. 2003a), its populations dynamics and reproductive biology have been studied (Reed, 1995a, 1995b; Aldana Aranda et al. 2003b; 2003c; 2003d; Delgado et al 2004; Castro et al. 2005). It is from this reproduction related histological work that the presence of a parasite in the digestive gland was detected. Apicomplexa parasites are of common occurrence in invertebrates and especially in Mollusks (Lester & Davis, 1981; Hillman et al, 1982; Perkins, 1988; Duszynski et al., 1999, 2004). The goal in this study was to identify the abundance of the different stages of the Apicomplexa like parasite from the digestive gland of *S. gigas* from different localities in the Caribbean region.

## MATERIAL AND METHODS

Samples for histological study of *S. gigas* were made at San Andres, Colombia; Alacranes, Cozumel and Chinchorro from Mexico; Tourmaline, Puerto Rico; Guadeloupe and San Berthelemy, French Antilles and Margarita Island from Venezuela (Fig. 1). They were sampled one or more times to ascertain the presence and abundance of the parasite, including some laboratory reared juveniles from Florida that were imported to Xcaret, Mexico. All conchs sampled were adult with a shell lip over 6 mm thick. A sample of 1 cm<sup>3</sup> of the spire, including digestive gland (DG) and gonad (G) was processed through standard histological techniques (Gabe, 1968). Sections were stained with a modified Goldner trichrome method, including Hematoxylin, acid fuchsin and phosphomolybdic orange G with Alcian blue. Digital images were taken with a Sony CCD-IRIS video-camera mounted on the Carl Zeiss microscope or a Nikon DXm 1200F digital camera mounted on the Nikon microscope. The quantification to establish an incidence of the parasite was done counting the total of every stage observed in two 40x fields per slide, ten slides of different organisms per month and locality were used.

## RESULTS

The different stages identified in the digestive gland of *S. gigas* were an Apicomplexa like, bottle shaped trophozoites, 25-30 µm long and 15-20 µm wide, with conic apical structure, it is always found within the cells (Pl. 1 a-b). The apical structure is implanted in the cellular wall (Pl. 1b). These trophozoites become cysts (gametocyst) with no connection to the host cell (Pl. 1c). Gametes 15 to 20 µm with two to four macrogametes 12 to 15 µm. (Pl. 1c) or microgametes 5-8 µm (Pl. 2b) and Oocyst with a single trophozoite (Pl. 1d), and Oocyst with four trophozoites (Baquero, et al. in press).

All sampled organisms presented one or more stages of the parasite in the digestive gland. Figure 2 presents the variation in abundance of the different stages of the parasite at the localities studied. The dominant stage of the

parasites varies with the site and in time, even though, the gametocysts are the dominant stage through the Caribbean the whole year. The lowest incidence was registered in the laboratory reared organisms from Xcaret with a total of 9 parasites per field, and the highest were registered at Puerto Rico with 85 parasites per field and San Andres with 82 parasites per field. The mean number of parasites at San Andres and Alacranes through the year was similar, 54 and 50 respectively (Fig. 3).

The different stages vary much in number through the year, with variations for trophozoites of 1 to 7 individuals per field, gametocysts from 3 to 47, gametes from 1 to 45 (Fig. 4). Gametes and oocysts are absent during several months but they can be very abundant, gametes are found in groups of over 30 with several groups per field, while oocysts were registered up to 25 per field. Oocysts start from the conjugation of a macro and microgamete (Duszynski, et al. 2004), originating a small cyst, easily distinguished from gametocysts that are originated from the encystment of a trophozoite, but as they mature and enlarge they can be mistaken with gametocyst.

The variation of stage abundance and chronology was investigated at Alacranes, Mexico and San Andres, Colombia where a year cycle was sampled. Relative abundance of the different stages between the two localities is no similarity in these localities. The dominant stage is the gametocyst, followed by gametes, except for December at Alacranes that the dominant stage is the trophozoites. The largest number of gametocysts was detected at Alacranes from August to November, while at San Andres it was from January to June.

## DISCUSSION


The dominant stage at any moment are generally the gametocysts, but their importance seem to vary with the time of the year, presenting different patterns at every locality, which may be influenced by the prevailing environ-




**Figure 1.** Samples for histological study of *Strombus gigas* in different localities from the Caribbean region.

mental conditions. This is clearly seen when Alacranes and San Andres populations were compared. But it is also possible that a variation in abundance is associated with the physiological stage of the conch, taking advantage of the renewal of the digestive cell.

The presence of cysts and spores of an intracellular parasitic organism in the digestive gland cells is constant, although it varies in intensity according to individuals. Their presence in the lumen of the digestive gland ducts is much more variable, probably according to the discharge



**Figure 2.** Variation in abundance of the different stages of the parasite at the localities studied.



**Figure 3.** Incidence of the Apicomplexa infecting the digestive gland of *S. gigas* in different localities in the Caribbean.

cycle of secretory cells. Gestala et al. (2002) described on the life cycle of *Aggregata* (Apicomplexa, Aggregatidae).

The constant presence of the different stages of the parasite implies that infection occurs all the year round. As maturing gametocysts and gamots were detected emerging in the same organism, it can be interpreted as a re-infection from the ingestion of gametocysts and oocysts, or that they complete the life cycle without leaving the host. The second hypothesis is more likely to be the regular process, as just in a few cases a total destruction of the digestive diverticula cells was detected, with a massive liberation of every present stage. The different stages identified throughout the Caribbean suggest that the parasite may be an Eimeriidae, which require of two hosts to complete their life cycle (Duszynski et al. 2004). But it is not rare that they complete the life cycle within the same host with the gametocysts expelled with the feces as a dispersion stages (Duszynski, et al. 1999). Apparently, the organisms found in the digestive gland of *S. gigas* belong to the second modality, as all the stages have been found within the digestive gland tissue. It is possible that we are dealing with more than one species, as suggested by an oocyst found at Guadalupe with four trophozoites within, compared to the frequently found one trophozoite oocyst at all other localities (Pl. 3).

The intracellular characteristic of the infection, the presence of an apical feeding structure of the trophozoite, and of sporozoite stages with macro and microgamonts, suggest this may be a Coccidia, however they are reported mostly for vertebrates. Pl. 4a and 4d could be spores of *Nematopsis* spp. No reference has been found of such large intracellular trophozoites, being similar only to extracellular parasites of the family Eimeridae, which is characterized for having more than one host (Bower, 2001)

The presence of such a generalized and intense infection and its coincidence with reduced reproductive activity raises several questions: What are the environmental factors inducing such an intense and generalized infection? What is the geographic extension of the disease and its impact on recruitment and long term health of impacted populations? Is it possible to revert to a healthy state? And

last but not least, what is the putative danger of transmission to natural predators and humans when the queen conchs are eaten raw?

### ACKNOWLEDGMENTS

Ichthyology laboratory of CINVESTAVIPN, Mérida for its support for histology processing. Thanks to Teresa Colas for support in histological process and digital images.

### LITERATURE CITED

- Aldana Aranda, D. Erick Baqueiro., Martínez, I., Ochoa, RI and brulé, T. 2001. Reproductive patterns of *Strombus gigas* from Alacranes reef versus Chinchorro bank of Mexico. *Proceeding Gulf and Caribb. Fish. Instit.* **54** :202-224p.
- Aldana Aranda D., E. Baqueiro Cardenas & S. Manzanilla Naim. 2003a. Mexican marine parks as a fishery management tool for the queen conch *Strombus gigas*. In: Aldana-Aranda, D. (Ed.) *El caracol Strombus gigas: conocimiento integral para su manejo sustentable en el Caribe, México*. España: CYTED. pp.101-108.
- Aldana-Aranda, D., E. R. Baqueiro-Cárdenas, I. Martínez-Morales, R. I. Ochoa & T. Brulé. 2003b. Reproductive Patterns of *Strombus gigas* from Alacranes Reef versus Chinchorro Bank of Mexico. *Gulf and Caribbean Fisheries Institute* **54**: 202-225.
- Aldana-Aranda, D., E. R. Baqueiro-Cárdenas, I. Martínez-Morales, A. Zetina-Zarate & T. Brulé. 2003c. A review of the reproductive patterns of gastropods mollusks from Mexico. *Bulletin of Marine Science* **73**: 629-641.
- Aldana-Aranda, D., E. R. Baqueiro-Cárdenas, I. Martínez-Morales, R. I. Ochoa & T. Brulé. 2003d. Gonad behavior during peak reproduction period of *Strombus gigas* from Banco Chinchorro. *Bulletin of Marine Science* **73**: 241-248.
- Appeldoorn, R. S., 1987 Considerations and preliminary calculations of maximum sustainable yield for the queen conch (*Strombus gigas*) resource of Puerto Rico and the U.S. Virgin Island. Rep. Caribb. Fish. Manage. Counc. Dec. 1987:1-20.
- Baqueiro Cárdenas, E., L. Frenkiel & D. Aldana Aranda. 2005. One more threat for the Queen conch *Strombus gigas*: coccidian (Apicomplexa) infection of *S. gigas* digestive gland (preliminary results). *Gulf and Caribbean Fisheries Institute*. Vol **58**:17-18
- Baqueiro Cárdenas, E., L. Frenkiel & D. Aldana Aranda. Apicomplexa infection of digestive gland *S. gigas* Queen conch *Strombus gigas*. *J. Shellfish. Research*, in press
- Castro, E., L. Frenkiel, E. Baqueiro & D. Aldana Aranda. 2005. Atypical reproductive cycle of the queen conch *Strombus gigas* (Mollusca: Gastropoda). *Gulf and Caribbean Fisheries Institute* **58** (in press).
- Delgado, G.A., C.T. Bartels, R.A. Glazer, N.J. Brown-

- Peterson & K.J. McCarthy. 2004. Translocation as a strategy to rehabilitate the queen conch (*Strombus gigas*) population in the Florida Keys. *Fishery Bulletin*. **102**: 278-288.
- Duszynski, D.W. S.J. Upton & I. Couch. 1999. Coccidia (*Eimeria* and *Isospora*) of Invertebrates. Department of Biology, University of New Mexico. <http://biology.unm.edu/biology/coccidia>
- Duszynski, D.W. S.J. Upton & I. Couch.. 2004. The Coccidian of the World. Department of Biology, University of New Mexico. <http://biology.unm.edu/biology/coccidia>
- Gabe, M.1968. Technique histologique Masson, Paris 1168p
- Gestala, C., A., S. Guerra Pascual & C. Azevedo. 2002. On the life cycle of *Aggregata eberthi* and observations on *Aggregata octopiana* (Apicomplexa, Aggregatidae) from Galicia (NE Atlantic). *European Journal of Protistology*. **37**: 427-435
- Hillman, R. E., N. J. Maciolek, J. I. Lahey & C. I. Belmore. 1982. Effects of haplosporidian parasite, *Haplosporidium* sp. on species of the molluscan woodborer *Teredo* in Barnegat Bay, New Jersey. *J. Invertebr. Pathol.* **40**: 307-319.
- Lester, R. J. F. & G.H., Davis. 1981. A new Perkinsus species (Apicomplexa, Perkinsea) from the abalone *Haliotis ruber*. *J. Invertebrate Pathol.* **37**:208-209.
- Perkins, O. F.1988. Parasite morphology, strategy and evolution. In: W. S. Fisher, editors. Disease Processes in Marine Bivalve mollusks. US: *American Fisheries Soc.* Special Publication **18**. pp. 93-111.
- Reed, S. E. 1995a. Reproductive anatomy and biology of the genus *Strombus* in the Caribbean: I. Males. *Journal of Shellfish Research* **14**: 325-330.
- Reed, S. E. 1995b. Reproductive anatomy and biology of the genus *Strombus* in the Caribbean: II. Females. *Journal of Shellfish Research* **14**: 331-336.
- Voltzow, J. 1994. Gastropoda: Prosobranchia. In: F.W. Harrison & A.J. Kohn, editors. Microscopic anatomy of invertebrates, Vol 5, Mollusca I. New York: John Wiley and Sons Ltd. pp. 111-252.

